




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C07C 401/00, A61K 31/59	A1	(11) International Publication Number: WO 90/09991 (43) International Publication Date: 7 September 1990 (07.09.90)
(21) International Application Number: PCT/DK90/00036 (22) International Filing Date: 13 February 1990 (13.02.90) (30) Priority data: 8904154.5 23 February 1989 (23.02.89) GB (71) Applicant (for all designated States except US): LEO PHARMACEUTICAL PRODUCTS LTD. A/S (LØVENS KEMISKE FABRIK PRODUKTIONS AKTIESELSKAB) [DK/DK]; Industriparken 55, DK-2750 Ballerup (DK). (72) Inventors; and (75) Inventors/Applicants (for US only): CALVERLEY, Martin, John [GB/DK]; Oktobervej 61, DK-2730 Herlev (DK). HANSEN, Kai [DK/DK]; Lidsøvej 45, DK-2730 Herlev (DK). BINDERUP, Lise [DK/DK]; Ludvig Hegners Allé 8A, DK-2630 Tåstrup (DK).		(74) Agent: KRISTENSEN, Rydahl, P.; Patent Dept., Leo Pharmaceutical Products, Industriparken 55, DK-2750 Ballerup (DK). (81) Designated States: AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (European patent), FI, FR (European patent), GB (European patent), HU, IT (European patent), JP, KR, LU (European patent), NL (European patent), NO, RO, SE (European patent), SU, US. Published <i>With international search report.</i>

(54) Title: NOVEL VITAMIN D ANALOGUES**(57) Abstract**

The present invention relates to compounds of formula (I), in which formula R stands for an alkyl group containing from 7 to 12 carbon atoms optionally substituted with a hydroxy group; and derivatives of the compounds of formula (I) in which one or more hydroxy groups have been transformed into -O-acyl or -O-glycosyl or phosphate ester groups, such masked groups being hydrolyzable *in vivo*, in pure form or in mixtures. The present compounds find use in both the human and veterinary practice in the treatment and prophylaxis of autoimmune diseases, including diabetes mellitus, hypertension, inflammatory diseases such as rheumatoid arthritis and asthma as well as diseases characterized by abnormal cell differentiation and/or cell proliferation, and/or imbalance in the immune system.

FOR THE PURPOSES OF INFORMATION ONLY


Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MR	Mauritania
BE	Belgium	GA	Gabon	MW	Malawi
BF	Burkina Faso	GB	United Kingdom	NL	Netherlands
BG	Bulgaria	HU	Hungary	NO	Norway
BJ	Benin	IT	Italy	RO	Romania
BR	Brazil	JP	Japan	SD	Sudan
CA	Canada	KP	Democratic People's Republic of Korea	SE	Sweden
CF	Central African Republic	KR	Republic of Korea	SN	Senegal
CG	Congo	LI	Liechtenstein	SU	Soviet Union
CH	Switzerland	LK	Sri Lanka	TD	Chad
CM	Cameroon	LU	Luxembourg	TG	Togo
DE	Germany, Federal Republic of	MC	Monaco	US	United States of America
DK	Denmark				

NOVEL VITAMIN D ANALOGUES


This invention relates to a hitherto unknown class of compounds which shows an immunomodulating effect as well as strong activity in inducing differentiation and inhibiting undesirable proliferation of certain cells, including cancer cells and skin cells, to pharmaceutical preparations containing these compounds, to dosage units of such preparations, and to their use in the treatment and prophylaxis of autoimmune diseases, including diabetes mellitus, hypertension, inflammatory diseases such as rheumatoid arthritis and asthma as well as diseases characterized by abnormal cell differentiation and/or cell proliferation, and/or imbalance in the immune system.

The compounds of the present invention are represented by the general formula I



in which formula R stands for an alkyl group containing from 4 to 12 carbon atoms optionally substituted with a hydroxy group.

Preferably R is a group of formula II



5 where n is an integer from 1 to 7; R¹ and R², which may be the same or different, stand for hydrogen, lower alkyl, lower cycloalkyl, or, taken together with the carbon atom (starred in formula II) bearing the group X, R¹ and R² can form a C₃-C₈ carbocyclic ring; X stands for hydrogen or
 10 hydroxy.

In the context of this invention, the expression "lower alkyl" indicates a straight or branched saturated or unsaturated carbon chain containing from 1 to 5 carbon atoms, and the expression "lower cyclo-alkyl" indicates a
 15 saturated or unsaturated C₃-C₇ carbocyclic ring.

As can be seen from formula I and II, depending on the meanings of R, X, R¹ and R², the compounds of the invention can comprise several diastereoisomeric forms (e.g. R or S configuration at the starred carbon atom). The in-
 20 vention covers all these diastereoisomers in pure form and also mixtures of diastereoisomers. In addition, derivatives of I in which one or more of the hydroxy groups are masked as groups which can be reconverted to hydroxy groups in vivo are also within the scope of the invention ("biore-
 25 versible derivatives or prodrugs of I").

The term "bioreversible derivatives or prodrugs of I" includes, but is not limited to, derivatives of the compounds of formula I in which one or more hydroxy groups have been transformed into -O-acyl or -O-glycosyl groups,
 30 or a phosphate ester, such masked groups being hydrolyzable in vivo.

Compounds of formula I in which R is not substituted with hydroxy are another type of prodrug. These compounds are relatively inactive in vitro, but are converted to
 35 active compounds of formula I by enzymatic hydroxylation after administration to the patient.

It has recently been shown that 1 α ,25-dihydroxy-vitamin D₃ (1,25(OH)₂D₃) influences the effects and/or

production of interleukins, indicating the potential use of this compound in the treatment of diseases characterized by a dysfunction of the immune system, e.g. autoimmune diseases and rejection of transplants. In addition, other conditions characterized by an abnormal interleukin-1 production, e.g. inflammatory diseases such as rheumatoid arthritis may be treated with $1,25(\text{OH})_2\text{D}_3$.

It has also been shown that $1,25(\text{OH})_2\text{D}_3$ is able to stimulate the differentiation of cells and inhibit excessive cell proliferation, and it has been suggested that this compound might be useful in the treatment of diseases characterized by abnormal cell proliferation and/or cell differentiation such as cancer and psoriasis.

Also, the use of $1,25(\text{OH})_2\text{D}_3$ for the treatment of hypertension and diabetes mellitus has been suggested.

However, the therapeutic possibilities in such indications of $1,25(\text{OH})_2\text{D}_3$ are severely limited by the well known potent effect of this hormone on calcium metabolism; elevated blood concentrations will rapidly give rise to hypercalcemia. Thus, this compound and its potent synthetic analogues are not completely satisfactory for use as drugs in the treatment of e.g. psoriasis, cancer or immune diseases which may require continuous administration of the drug in relatively high doses.

A number of oxa-analogues of vitamin D_3 are known. $1\alpha,25$ -dihydroxy-20-oxa-21-norvitamin D_3 and 1α -hydroxy-20-oxa-21-norvitamin D_3 are described in N. Kubodera et al, Chem. Pharm. Bull., 34, 2286 (1986), $1\alpha,25$ -dihydroxy-22-oxavitamin D_3 and 25-hydroxy-22-oxavitamin D_3 are described in E. Murayama et al, Chem. Pharm. Bull., 34, 4410 (1986), J. Abe et al, FEBS LETTER, 226, 58 (1987) and European Patent Application, publication number 184 112, and $1\alpha,25$ -dihydroxy-23-oxavitamin D_3 is described in European Patent Application, publication number 78704.

In vitro experiments indicate that some of these compounds may have advantages over $1,25(\text{OH})_2\text{D}_3$. Thus $1\alpha,25$ -dihydroxy-22-oxavitamin D_3 has only one 14th as much affinity as $1\alpha,25(\text{OH})_2\text{D}_3$ for the chick intestinal cytosolic

receptor, a weaker affinity than $1,25(\text{OH})_2\text{D}_3$ for the receptor in a human myeloid leukemia cell line (HL-60), and a high activity as inducer of differentiation in HL-60 cells.

- 5 In contrast to the compounds of the present invention the above mentioned 22-oxa-compounds have the S-configuration in the 20-position.

The usefulness of a vitamin D analogue in the above mentioned indications is dependent not only upon a favourable ratio of binding affinity to relevant receptors compared to the intestinal receptor, but also upon the fate of the compound in the organism.

10 It has now been found that the compounds of the present invention show favourable selectivity with respect to receptor binding and at the same time show high bioavailability as well as chemical and metabolic stability.

The selectivity of the compounds is illustrated by the fact that while they have high affinities for the receptor in tumour cells (similar to or much better than that of $1,25(\text{OH})_2\text{D}_3$) and the concentration needed to induce cell differentiation in a human monocytic tumour cell line is the same as or considerably lower than that needed of $1,25(\text{OH})_2\text{D}_3$ to give the same effect, their binding affinity for the intestinal receptor is lower than that of $1,25(\text{OH})_2\text{D}_3$. In vivo in rats the compounds are less active than $1,25(\text{OH})_2\text{D}_3$ in inducing hypercalciuria and hypercalcemia.

This renders the compounds of the invention especially suited for both local and systemic treatment and prophylaxis of human and veterinary disorders which are characterized by abnormal cell proliferation and/or cell differentiation, such as certain dermatological disorders including psoriasis and certain cancer forms, e.g. leukemia and myelofibrosis, and diseases characterized by an imbalance in the immune system, e.g. autoimmune diseases, or AIDS, and to obtain desired immunosuppression as in transplantation procedures, as well as treatment of acne, diabetes mellitus and hypertension and inflammatory diseases, such

as rheumatoid arthritis and asthma. As the compounds of this invention may promote the differentiation of the hair follicle cells, these compounds may be used in the treatment of alopecia.

5 The compounds of formula I may conveniently be prepared from the vitamin D-derivative 1 (or its 20R isomer) (Tetrahedron, 43, 4609 (1987)) by the routes outlined in Scheme 1. Oxidation of 1 for example using the van Rhee procedure (Tetrahedron Letters, 1969, 985) gives the ketone
10 2, which is reduced to the 20R-alcohol 3. When a suitable chiral reducing agent is used 3 may be prepared with very high stereoselectivity, but 3 is conveniently prepared by NaBH_4 reduction of 2 and separating the minor amount of corresponding 20S-alcohol chromatographically. O-Alkylation
15 of 3 to give III is achieved by treatment under basic conditions with a side chain building block of general formula Z-R^3 , in which Z is a leaving group such as a halogen (Cl, Br or I) or p-toluenesulphonyloxy or methanesulphonyloxy, and R^3 is R (of formula I) or optionally a
20 radical which can be converted to R at any convenient later stage (or over several stages). Thus R^3 in compounds III, IV, V and VI does not necessarily have the same meaning along a particular synthetic sequence. The conversion of R^3 to R may well involve several steps and possibly involve a
25 temporary protection of the sensitive triene system of the molecule. Apart from any necessary modification within the side chain (R^3), the conversion of III to I involves a photoisomerisation step and a desilylation step, analogous to the steps used in the last stages of the synthesis of
30 other vitamin D analogues (see European patent No. 0 227 826).

 The side chain building blocks, R^3Z , are either known compounds (several are described in international patent application PCT/DK89/00079) or may be prepared analogously
35 to those described in PCT/DK89/00079. The R^3 is typically identical with formula II in which X is a protected OH group, e.g. tetrahydropyranyloxy or trialkylsilyloxy. (Any such THP ethers R^3Z , which are not described in

PCT/DK89/00079, are readily prepared from the corresponding alcohol).

The following standard abbreviations are used throughout this disclosure: Me = methyl; Et = ethyl; Prⁿ = n-propyl; Prⁱ = isopropyl; Bu^t = tert-butyl; THP = tetrahydro-4H-pyran-2-yl; THF = tetrahydrofuran; Ts = p-toluene-sulphonyl; TBA = tetra-(n-butyl)-ammonium.

10

15

20

25

30

35

Notes to Scheme 1

- a) Oxidation e.g. with O_2 with $Cu(AcO)_2$, 2,2'-bipyridyl and 1,4-diazabicyclo[2,2,2]octane as catalyst.
- 5 b) Reduction (e.g. with $NaBH_4$).
- c) Alkylation with the side chain fragment R^3-Z in the presence of base (e.g. KOH , $KOBu^t$ or KH , with or without catalyst (e.g. 18-Crown-6) in solvent, e.g. THF.
- 10 d) Optional functional group modification in the side chain.
- e) Isomerisation with $h\nu$ - triplet sensitizer, e.g. anthracene.
- f) Deprotection with TBA^+F^- or HF .

15 It should be noted that although the shown intermediates may have hydroxyl groups protected as tert-butyl-dimethylsilyl ethers, the scope of the invention does not exclude the use of alternative hydroxyl protecting groups well known in the art (such as those described in T.W. Greene, "Protective groups in organic synthesis", Wiley, New York, 1981), together with alternative reactions for deprotection.

The present compounds are intended for use in pharmaceutical compositions which are useful in the treatment of human and veterinary disorders as described above.

The amount required of a compound of formula I (hereinafter referred to as the active ingredient) for therapeutic effect will, of course, vary both with the particular compound, the route of administration and the mammal under treatment. The compounds of the invention can be administered by the parenteral, intra-articular, enteral or topical routes. They are well absorbed when given enterally and this is the preferred route of administration in the treatment of systemic disorders. In the treatment of dermatological disorders like psoriasis, topical or enteral forms are preferred.

In the treatment of respiratory diseases like asthma an aerosol is preferred.

While it is possible for an active ingredient to be administered alone as the raw chemical, it is preferable to present it as a pharmaceutical formulation. Conveniently, the active ingredient comprises from 1 ppm to 0.1% by weight of the formulation.

By the term "dosage unit" is meant a unitary, i.e. a single dose which is capable of being administered to a patient, and which may be readily handled and packed, remaining as a physically and chemically stable unit dose comprising either the active material as such or a mixture of it with solid or liquid pharmaceutical diluents or carriers.

The formulations, both for veterinary and for human medical use, of the present invention comprise an active ingredient in association with a pharmaceutically acceptable carrier therefore and optionally other therapeutic ingredient(s). The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulations and not deleterious to the recipient thereof.

The formulations include e.g. those in a form suitable for oral, rectal, parenteral (including subcutaneous, intramuscular and intravenous), intra-articular and topical administration.

The formulations may conveniently be presented in dosage unit form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing the active ingredient into association with the carrier which constitutes one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing the active ingredient into association with a liquid carrier or a finely divided solid carrier or both, and then, if necessary, shaping the product into the desired formulation.

Formulations of the present invention suitable for oral administration may be in the form of discrete units as capsules, sachets, tablets or lozenges, each containing a predetermined amount of the active ingredient; in the form of a powder or granules; in the form of a solution or a

suspension in an aqueous liquid or non-aqueous liquid; or in the form of an oil-in-water emulsion or a water-in-oil emulsion. The active ingredient may also be administered in the form of a bolus, electuary or paste.

5 A tablet may be made by compressing or moulding the active ingredient optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing, in a suitable machine, the active ingredient in a free-flowing form such as a powder or granules, optionally
10 mixed by a binder, lubricant, inert diluent, surface active or dispersing agent. Moulded tablets may be made by moulding, in a suitable machine, a mixture of the powdered active ingredient and suitable carrier moistened with an inert liquid diluent.

15 Formulations for rectal administration may be in the form of a suppository incorporating the active ingredient and a carrier such as cocoa butter, or in the form of an enema.

 Formulations suitable for parenteral administration
20 conveniently comprise a sterile oily or aqueous preparation of the active ingredient which is preferably isotonic with the blood of the recipient.

 Formulations suitable for intra-articular administration may be in the form of a sterile aqueous preparation of
25 the active ingredient which may be in microcrystalline form, for example, in the form of an aqueous microcrystalline suspension. Liposomal formulations or biodegradable polymer systems may also be used to present the active ingredient for both intra-articular and ophthalmic administration.
30

 Formulations suitable for topical administration include liquid or semi-liquid preparations such as liniments, lotions, applicants, oil-in-water or water-in-oil emulsions such as creams, ointments or pastes; or solutions
35 or suspensions such as drops.

 For asthma treatment inhalation of powder, self-propelling or spray formulations, dispensed with a spray can, a nebulizer or an atomizer can be used. The formu-

lations, when dispensed, preferably have a particle size in the range of 10 to 100 μ .

Such formulations are most preferably in the form of a finely comminuted powder for pulmonary administration from a powder inhalation device or self-propelling powder-dispensing formulations. In the case of self-propelling solution and spray formulations, the effect may be achieved either by choice of a valve having the desired spray characteristics (i.e. being capable of producing a spray having the desired particle size) or by incorporating the active ingredient as a suspended powder in controlled particle size. These self-propelling formulations may be either powder-dispensing formulations or formulations dispensing the active ingredient as droplets of a solution or suspension.

Self-propelling powder-dispensing formulations preferably comprise dispersed particles of solid active ingredients, and a liquid propellant having a boiling point below 18°C at atmospheric pressure. The liquid propellant may be any propellant known to be suitable for medicinal administration and may comprise one or more C₁-C₆-alkyl hydrocarbons or halogenated C₁-C₆-alkyl hydrocarbons or mixtures thereof; chlorinated and flourinated C₁-C₆-alkyl hydrocarbons are especially preferred. Generally, the propellant constitutes 45 to 99.9% w/w of the formulation whilst the active ingredient constitutes 1 ppm to 0.1% w/w, of the formulation.

In addition to the aforementioned ingredients, the formulations of this invention may include one or more additional ingredients such as diluents, buffers, flavouring agents, binders, surface active agents, thickeners, lubricants, preservatives, e.g. methyl hydroxybenzoate (including anti-oxidants), emulsifying agents and the like.

The compositions may further contain other therapeutically active compounds usually applied in the treatment of the above mentioned pathological conditions.

The present invention further concerns a method for treating patients suffering from one of the above patholog-

ical conditions, said method consisting of administering to a patient in need of treatment an effective amount of one or more compounds of formula I, alone or in combination with one or more other therapeutically active compounds usually applied in the treatment of said pathological conditions. The treatment with the present compounds and/or with further therapeutically active compounds may be simultaneous or with intervals.

In the treatment of systemic disorders daily doses of from 0.1-100 μg , preferably from 0.2-25 μg , of a compound of formula I are administered. In the topical treatment of dermatological disorders, ointments, creams or lotions containing from 0.1-500 $\mu\text{g/g}$, and preferably from 1-100 $\mu\text{g/g}$, of a compound of formula I are administered. The oral compositions are formulated, preferably as tablets, capsules, or drops, containing from 0.05-50 μg , preferably from 0.1-25 μg , of a compound of formula I, per dosage unit.

The invention will now be further described in the following non-limiting Preparations and Examples:

Preparations and Examples

General

The exemplified compounds I are listed in Table 1. The intermediates of Scheme I referred to in the Preparations are to be identified by numbers with the corresponding formulae in Table 2.

For nuclear magnetic resonance spectra (300 MHz) chemical shift values (δ) are quoted for deuteriochloroform solutions relative to internal tetramethylsilane ($\delta = 0$) or chloroform ($\delta = 7.25$). The value for a multiplet, either defined (doublet (d), triplet (t), quartet (q)) or not (m) at the approximate mid point is given unless a range is quoted (s = singlet, b = broad). Coupling constants (J) are given in Hertz, and are sometimes approximated to the nearest unit.

Ether is diethyl ether, and was dried over sodium. THF was dried over sodium-benzophenone. Petroleum ether

refers to the pentane fraction. Reactions were run at room temperature unless otherwise noted. The work-up procedure referred to involves dilution with the specified solvent (otherwise the organic reaction solvent), extraction with water and then brine, drying over anhydrous MgSO_4 , and concentration in vacuo to give a residue.

Table 1: Exemplified Compounds I (R given in formula II)

	Compound Number	Example Number	Formula (II)			
			n	R^1	R^2	X
10	101	2	1	H	Pr^1	OH
	102	3	2	Me	Me	OH
15	103	12	2	$-(\text{CH}_2)_5-$		OH
	104	11	3	H	H	OH
	105	4	3	Me	Me	OH
	106	5	3	Et	Et	OH
	107	9	3	Pr	Pr	OH
20	108	10	4	Me	Me	H
	109	1	4	Me	Me	OH
	110	6	4	Et	Et	OH
	111	7	5	H	H	OH
	112	8	5	Me	Me	OH
25	113	13	6	Me	Me	OH

Table 2:

5	Com- pound Number	Prepar- ation Number	Formula	
			Type (See Scheme 1)	R ³
	4	11	III	-CH ₂ -CH=CMe ₂
	5	33	III	-(CH ₂) ₄ -CHMe ₂
	6	9	III	-CH ₂ -CH[OSi(Me ₂)Bu ^t]CHMe ₂
10	7	12	III	-(CH ₂) ₂ -C(OH)Me ₂
	8	37	III	-(CH ₂) ₂ -C(OH)-(CH ₂) ₄ -CH ₂
	9	31	III	-(CH ₂) ₄ -OSi(Me ₂)Bu ^t
	10	15	III	-(CH ₂) ₃ -C(O-THP)Me ₂
	11	14	III	-(CH ₂) ₃ -C(O-THP)Et ₂
15	12	26	III	-(CH ₂) ₃ -C(OSiMe ₃)Et ₂
	13	32	III	-(CH ₂) ₃ -C(OSiMe ₃)Pr ⁿ ₂
	14	3	III	-(CH ₂) ₄ -C(O-THP)Me ₂
	15	18	III	-(CH ₂) ₄ -C(OSiMe ₃)Me ₂
	16	19	III	-(CH ₂) ₄ -C(OSiMe ₃)Et ₂
20	17	22	III	-(CH ₂) ₆ -OSiMe ₂)Bu ^t
	18	28	III	-(CH ₂) ₅ -C(OSiMe ₃)Me ₂
	19	39	III	-(CH ₂) ₆ -C(OSiMe ₃)Me ₂
	20	35	IV	-(CH ₂) ₄ -CHMe ₂
	21	13	IV	-(CH ₂) ₂ -C(OH)Me ₂
25	22	38	IV	-(CH ₂) ₂ -C(OH)-(CH ₂) ₄ -CH ₂
	23	36	IV	-(CH ₂) ₄ -OSi(Me ₂)Bu ^t
	24	16	IV	-(CH ₂) ₃ -C(O-THP)Me ₂

Table 2 (continued):

5	Com- pound Number	Prepar- ation Number	Formula	
			Type (See Scheme 1)	R ³
	25	17	IV	$-(CH_2)_3-C(O-THP)Et_2$
	26	27	IV	$-(CH_2)_3-C(OSiMe_3)Et_2$
	27	34	IV	$-(CH_2)_3-C(OSiMe_3)Pr^n_2$
10	28	25	IV	$-(CH_2)_4-C(OSiMe_3)Me_2$
	29	24	IV	$-(CH_2)_4-C(OSiMe_3)Et_2$
	30	23	IV	$-(CH_2)_6-OSi(Me_2)Bu^t$
	31	29	IV	$-(CH_2)_5-C(OSiMe_3)Me_2$
	32	40	IV	$-(CH_2)_6-C(OSiMe_3)Me_2$
15	33	4	V	$-(CH_2)_4-C(O-THP)Me_2$
	34	10	VI	$-CH_2-CH(OH)CHMe_2$

Preparation 1: Compound 2

20 To a solution of 1(S),3(R)-bis-(tert-butyldimethyl-
silyloxy)-20(S)-formyl-9,10-secopregna-5(E),(7E),10(19)-
-triene (3.44 g, 6 mmol) (1) in N,N-dimethylformamide (150
ml), 1,4-diazabicyclo[2.2.2]octane (600 mg, 5.3 mmol),
cupric acetate, monohydrate (90 mg, 0.45 mmol) and
25 2,2'-bipyridyl (72 mg, 0.45 mmol) were added. Air was
bubbled through the well stirred solution for 6 days at
40°C.

The reaction mixture was diluted with ethyl acetate
(500 ml), extracted with water (2 x 100 ml) and saturated
30 aqueous sodium chloride (3 x 50 ml) and dried over MgSO₄.
Ethyl acetate was evaporated off, and the solid residue was
purified by chromatography (silica gel, 10% ether in petro-

leum ther as eluant) to give th title compound.

NMR: δ = 0.037 (s, 3H), 0.043 (s, 3H), 0.056 (s, 6H),
0.49 (s, 3H), 0.84 (s, 9H), 0.89 (s, 9H), 1.5-2.30 (m,
13H), 2.13 (s, 3H), 2.55 (dd, 1H), 2.70 (t, 1H), 2.89 (bd,
5 1H), 4.21 (m, 1H), 4.52 (m, 1H), 4.94 (m, 1H), 4.98 (m,
1H), 5.83 (d, 1H), 6.43 (d, 1H) ppm.

Preparation 2: Compound 3 and its 20S-isomer

Compound 2 (Prep. 1) (3.10 g, 5.5 mmol) was dissolved
10 in tetrahydrofuran (140 ml) and sodium borohydride (0.35 g,
3.3 mmol) was added. Methanol was then added dropwise over
15 minutes. The reaction blend was stirred for 20 minutes,
then diluted with ethyl acetate (560 ml). The solution was
extracted with water (5 x 150 ml) and saturated aqueous
15 sodium chloride (150 ml), dried over MgSO_4 and evaporated
to give a colourless oil. The oily residue was purified by
chromatography (silica gel, 15% ethyl acetate in petroleum
ether as eluant) and crystallization from methanol to give
3.

20 NMR: δ = 0.05 (m, 12H), 0.62 (s, 3H), 0.86 (s, 9H),
0.89 (s, 9H), 1.10-2.10 (m, 14H), 1.15 (d, 3H), 2.30 (bd,
1H), 2.53 (dd, 1H), 2.89 (m, 1H), 2.89 (m, 1H), 3.71 (m,
1H), 4.21 (m, 1H), 4.52 (m, 1H), 4.93 (m, 1H), 4.98 (m,
1H), 5.81 (d, 1H), 6.45 (d, 1H) ppm.

25 The fractions containing the more polar 20S-isomer
were evaporated to give a colourless residue which was
crystallized from methanol:

NMR, δ = 0.052 (bd, 12H), 0.54 (s, 3H), 0.85 (s, 9H),
0.89 (s, 9H), 1.22 (d, 3H), 1.20-2.10 (m, 14H), 2.30 (bd,
30 1H), 2.55 (dd, 1H), 2.87 (m, 1H), 3.72 (m, 1H), 4.21 (m,
1H), 4.52 (m, 1H), 4.94 (bs, 1H), 4.98 (m, 1H), 5.82 (d,
1H), 6.44 (d, 1H) ppm.

Preparation 3: Compound 14 (R^3 = 5-(tetrahydro-4-H-
-pyran-2-yloxy)-5-methyl-1-hexyl)

35 To a solution of compound 3 (561 mg, 1 mmol) in dry
tetrahydrofuran (10 ml) were added potassium hydroxide
(0.70 g, 10 mmol), 18-Crown-6 (40 mg) and 2-(6-bromo-2-

-methyl-2-hexyloxy)-tetrahydro-4H-pyran (Preparation 5a) (2.7 g, 10 mmol) were added. The mixture was stirred vigorously over the week-end, the reaction mixture was filtered, and the filtrate was evaporated in vacuo.

5 The residue was purified by chromatography (silica gel, 10% ether in petroleum ether as eluant) to give 14 as a colourless oil.

NMR, δ = 0.054 (m, 12H), 0.54 (s, 3H), 0.86 (s, 9H), 0.88 (s, 9H), 1.07 (d, J=6, 3H), 1.17 (s, 3H), 1.19 (s, 10 3H), 1.15-1.95 (m, 23H), 2.02 (t, 1H), 2.20 (bd, 1H), 2.30 (bd, 1H), 2.53 (dd, 1H), 2.85 (m, 1H), 3.10-3.30 (m, 2H), 3.40 (m, 1H), 3.55 (m, 1H), 3.93 (m, 1H), 4.20 (m, 1H), 4.51 (m, 1H), 4.69 (m, 1H), 4.93 (m, 1H), 4.98 (m, 1H), 5.79 (d, J=11, 1H), 6.45 (d, J=H, 1H) ppm.

15

Preparation 4: 1(S),3(R)-Dihydroxy-20(R)-[5'-(tetrahydro-4H-pyran-2"-yloxy)-5'-methyl-1'-hexyloxy]-9,10-seco-pregna-5(Z), 7(E),10(19)-triene (Compound 33)

20 A solution of compound 14 (400 mg, 0.5 mmol) anthracene (200 mg, 1.1 mmol) and triethylamine (1 drop) in dichloromethane (15 ml) under nitrogen in a Pyrex flask was irradiated with light from a high pressure ultra-violet lamp, type TQ 150Z2 (Hanau), at room temperature for 30 25 minutes. The solution was filtered and concentrated in vacuo to give the crude intermediate (Compound IV, Scheme 1, R^3 = 5-(tetrahydro-4-H-pyran-2-yloxy)-5-methyl-1-hexyl). This was dissolved in tetrahydrofuran (THF) (15 ml) and tetra-n-butylammonium fluoride trihydrate (1.05 g, 3.3 30 mmol) was added. The solution was heated at 60°C under an atmosphere of nitrogen for 1 hour. After cooling, the reaction mixture was partitioned between ethyl acetate (50 ml) and saturated aqueous sodium hydrogen carbonate (10 ml). The organic layer was washed with water (10 ml), dried and 30 concentrated. The residue was purified by chromatography (100 g, silica gel, 50% ethyl acetate in petroleum ether as eluant) to give the desired compound.

NMR, δ = 0.56 (s, 3H), 1.07 (d, 3H), 1.18 (s, 3H),

1.20 (s, 3H), 1.1-2.05 (m, 24H), 2.17 (bd, 1H), 2.30 (dd, 1H), 2.57 (dd, 1H), 2.81 (m, 1H), 3.10-3.30 (m, 2H), 3.42 (m, 1H), 3.56 (m, 1H), 3.93 (m, 1H), 4.22 (m, 3H), 4.41 (m, 1H), 4.70 (m, 1H), 5.00 (bs, 1H), 5.33 (bs, 1H), 5.99 (d, 1H), 6.39 (d, 1H) ppm.

Preparation 5a: 2-(6-Bromo-2-methyl-2-hexyloxy)-
-tetrahydro-4H-pyran

To a stirred, ice-cooled solution of ethyl 5-bromopentanoate (18.7 ml) in dried ether (100 ml) was added dropwise over 1 hour a filtered solution of Grignard reagent, prepared from magnesium (10 g) and methyl iodide (25 ml) in dried ether (200 ml). After a further 30 minutes on the ice-bath, the reaction mixture was allowed to warm to room temperature over 30 minutes before being poured onto a stirred, ice-cooled solution of ammonium chloride (30 g) in water (200 ml). After the vigorous reaction had subsided, the ether layer was separated, and the aqueous layer was extracted with more ether. The combined ether layers were washed consecutively with water and saturated aqueous sodium chloride, dried, and concentrated in vacuo to give the crude intermediate (6-bromo-2-methyl-2-hexanol) as a pale yellow oil. This was dissolved in dichloromethane (100 ml), then 3,4-dihydro-2H-pyran (8.9 ml) and pyridinium p-toluenesulfonate (0.8 g) were added at room temperature. After 1 hour, the reaction solution was diluted with ether (250 ml) and extracted consecutively with saturated aqueous sodium hydrogen carbonate (150 ml), water (100 ml) and saturated aqueous sodium chloride (100 ml). After drying and removal of the solvent in vacuo, the product was purified by chromatography (150 g silica gel, 10% ether in petroleum ether as eluant) to give the desired compound as a colourless oil.

NMR: δ = 1.20 (s, 3H), 1.22 (s, 3H), 1.40-1.95 (m, 12 H), 3.42 (t, 2H), 3.94 (m, 1H), 3.45 (m, 1H), 4.72 (m, 1H) ppm.

Preparation 5b: 2-(6-Bromo-3-ethyl-2-hexyloxy)-
-tetrahydro-4H-pyran

Using a procedure analogous to that of Preparation 5a, the title compound was prepared from ethyl 4-bromobutan-
5 oate, and the Grignard reagent derived from ethyl iodide.
NMR in agreement with structure.

Preparation 6: 5-Hydroxy-2,2,6-trimethyl-3(E)-
-heptene

10 To a solution of diethyl isobutyrylmethylphosphonate (22 g), tetrabutylammonium bromide (4 g) and pivaldehyde (13 ml) in dichloromethane (340 ml) was added 4 N aqueous sodium hydroxide solution (140 ml). The mixture was stirred overnight, and, after dilution with water, the organic
15 phase was worked up. The intermediate 5-oxo-2,2,6-trimethyl-3(E)-heptane was then isolated by distillation (b.p. 45-48°C/0.1 mBar). A stirred, ice-cooled solution of this (5 g) in a 0.4 M solution cerium III chloride in methanol (90 ml) was treated portionwise with sodium borohydride
20 (1.4 g). After 10 minutes, the mixture was worked-up (ethyl acetate) to give the title compound as an oil.

NMR: δ = 0.87 (d, 3H, J = 6.8), 1.02 (s, 9H), 1.50 (bs, 1H), 1.70 (m, 1H), 3.77 (bt, 1H), 5.36 (dd, 1H, J = 7.4 and 15.7), 5.65 (dd, 1H, J = 15.7 and 0.8).

25 NOTE: This racemic compound was resolved using Sharpless' kinetic resolution procedure (J. Amer. Chem. Soc. 1981, 103, 6237) to give either the S-form (using (-)-di-isopropyl tartrate) or the R-form (using (+)-di-isopropyl tartrate). These resolved forms can be used as starting
30 material for the following steps in the sequence here described for the conversion of the racemate to the side chain building block and then the target compound of Example 2.

Preparation 7: 3-Methyl-2-(tert-butyldimethylsilyl-
35 oxy)-butanal

A solution of 5-hydroxy-2,2,6-trimethyl-3(E)-heptene (Preparation 6) (4.5 g), imidazole (5 g) and tert-butyldimethylsilyl chloride (5 g) in dimethylformamide (50 ml) was

stirred for 1 hour. Work-up (ether) and distillation gave the intermediate 5-(tert-butyldimethylsilyloxy)-2,2,6-trimethyl-3(E)-heptene as an oil. (b.p. 65-69°C/0.03 mBar). A solution of this (7 g) in methanol (100 ml) and dichloromethane (320 ml) at -70°C was treated with ozonised oxygen until the reaction was judged to be complete (tlc analysis) (40 minutes), whereupon triphenylphosphine (9 g) was added and the reaction mixture allowed to warm to room temperature. Work-up (dichloromethane) and distillation gave the title compound as an oil, b.p. 45-48°C/1 mBar.

NMR δ = 0.04 (s, 6H), 0.90 (d, 3H), 0.92 (s, 9H), 0.95 (d, 3H), 2.01 (m, 1H), 3.70 (dd, 1H, J = 4.8 and 2.1), 9.58 (d, 1H, J = 2.1).

15 Preparation 8: 3-Methyl-2-(tert-butyldimethylsilyloxy)-1-(trifluoromethanesulfonyl)-oxy)butane

A stirred, ice-cooled solution of 3-methyl-2-(tert-butyldimethylsilyloxy)butanal (Preparation 7) (0.5 g) in THF (4 ml) and ethanol (8 ml) was treated with sodium borohydride (0.1 g). After 20 minutes, the reaction mixture was worked-up (ethyl acetate) to give the intermediate 3-methyl-2-(tert-butyldimethylsilyloxy)-1-butanol as an oil. This was dissolved in dichloromethane (5 ml), cooled to 0°C, and treated with pyridine (0.5 ml) and trifluoromethansulphonic anhydride (0.5 ml). After stirring for 1 hour, the reaction mixture was worked-up (ether) to give the title compound as an oil.

NMR: δ = 0.07 (s, 3H), 0.08 (s, 3H), 0.90 (s, 9H), 0.90 (d, 3H), 0.93 (d, 3H), 1.84 (m, 1H), 3.75 (m, 1H), 4.34 (dd, 1H, J = 9.9 and 6.8), 4.43 (d, 1H, J = 9.9 and 3.8).

Preparation 9: 1(S),3(R)-Bis(tert-butyldimethylsilyloxy)-20(R)-(3'-methyl-2'-tert-butyldimethylsilyloxy-1'-butoxy)-9,10-secopregna-5(E),7(E),10(19)-triene (Compound 6)

A stirred solution of Compound 3 (0.24 g), 18-Crown-6 (40 mg) and potassium tert-butoxide (0.15 g) in dry THF (4 ml) was treated with 3-methyl-2-tert-butyldimethylsilyloxy-1-trifluoromethanesulphonyloxybutane (Preparation 8) (0.3 g). After 15 minutes, the reaction mixture was worked-up (ether), and the residue purified by chromatography (silica gel, 2% ether in petroleum ether as eluant to give the title compound as an approx. equal mixture of diastereoisomers (epimers as position 2').

NMR δ = 0.0-0.12 (m, 18H), 0.53 and 0.54 (2s, 3H), 0.60-2.65 (m, 52H), 2.87 (m, 1H), 3.17 (m, 1H), 3.23 (m, 1H), 3.44 (m, 1H), 3.55 (m, 1H), 4.21 (m, 1H), 4.53 (m, 1H), 4.93 (m, 1H), 4.98 (m, 1H), 5.80 (d, 1H, J = 11.4), 6.46 (d, 1H, J = 11.4).

Preparation 10: 1(S),3(R)-Dihydroxy-20(R)-(2'-hydroxy-3'-methyl-1'-butoxy)-9,10-secopregna-5(E),7(E),10(19)-triene (Compound 34)

A stirred solution of 1(S),3(R)-bis(tert-butyldimethylsilyloxy)-20(R)-(3'-methyl-2'-tert-butyldimethylsilyloxy-1'-butoxy)-9,10-secopregna-5(E),7(E),10(19)-triene (Compound 6) (0.2 g) and tetrabutylammonium fluoride (0.7 g) in THF (5 ml) was heated at 60°C under nitrogen for 1 hour. After cooling, the reaction mixture was worked-up (ethyl acetate). Purification by chromatography (silica gel; ethyl acetate as eluant) gave the title compound.

NMR: δ = 0.58 and 0.60 (2s, 3H), 0.92 (d, 3H, J = 6.9), 0.98 (d, 3H, J = 6.9), 1.05-2.70 (m, 20H), 2.86 (m, 2H), 3.13-3.63 (m, 5H), 4.22 (m, 1H), 4.48 (m, 1H), 4.97 (m, 1H), 5.12 (m, 1H), 5.87 (d, 1H, J = 11.4), 6.57 (d, 1H, J = 11.4).

Preparation 11: 1(S),3(R)-Bis-[tert-butyl(dimethylsilyl)oxy]-20(R)-(3-methylbut-2-en-1-yloxy)-9,10-secopregna-5(E),7(E),10(19)-triene (Compound 4)

5 To a solution of compound 3 (0.61 g) in dry THF (10 ml) were added powdered potassium hydroxide (1.2 g), 18-Crown-6 (80 mg) and 3,3-dimethylallyl bromide (2.2 g). After stirring at room temperature for 24 hours, the mixture was partitioned between ether and water. The ether
10 layer was washed with brine, dried and concentrated in vacuo to give an oil. Purification by chromatography (silica gel; 2% to 5% ether in petroleum ether as eluent) followed by crystallization from methanol gave 4 as needles.

15 NMR: δ = 0.05 (bs, 12H), 0.55 (s, 3H), 0.86 (s, 9H), 0.89 (s, 9H), 1.10 (d, 3H), 1.65 (m, 3H), 1.72 (m, 3H), 1.05-1.82 (m, 10H), 1.90 (m, 1H), 2.03 (bt, 1H), 2.14 (m, 1H), 2.30 (m, 1H), 2.54 (dd, 1H), 2.87 (m, 1H), 3.30 (m, 1H), 3.78 (m, 1H), 4.06 (m, 1H), 4.21 (m, 1H), 4.52 (m,
20 1H), 4.93 (m, 1H), 4.98 (m, 1H), 5.33 (m, 1H), 5.80 (d, 1H, J = 11.5), 6.46 (d, 1H, J = 11.5).

Preparation 12: 1(S),3(R)-Bis-[tert-butyl(dimethylsilyl)oxy]-20(R)-(3-hydroxy-3-methyl-1-butoxy-9,10-secopregna-5(E),7(E),10(19)-triene (Compound 7)

25

NB: This preparation illustrates the protection of the triene system of III as an SO₂-adduct to allow efficient functional group modification in the side chain.

30

A solution of compound 4 (100 mg) in a few drops of ether was treated at -10°C with liquid sulphur dioxide (3 ml). The stirred mixture was allowed to warm spontaneously under a slow stream of nitrogen, and after 30 minutes the residual volatile material was removed on the rotary
35 evaporator. The residue was dissolved in THF (2 ml) and treated with a mixture prepared by adding THF (1 ml) to a solution of mercury II acetate (100 mg) in water (1 ml). The reaction mixture was stirred at 5°C for 18 hours and

then treated with 3N NaOH (3 ml) followed by a solution of NaBH₄ (0.05 g) in 3N NaOH (2 ml). Ethyl acetate was added and the mixture filtered through celite. The organic layer was washed with brine, dried and concentrated in vacuo to give a gum. This was dissolved/suspended in 96% ethanol (4 ml) together with sodium bicarbonate (0.2 g) and the stirred mixture was heated under reflux under nitrogen for 80 minutes. After cooling, the ethyl acetate was added and the mixture was extracted with water. The organic layer was washed with water. The organic layer was washed with brine, dried and concentrated in vacuo to give a residue. Purification by chromatography (silica gel, 5% to 30% ether in petroleum ether as eluent) gave 7.

NMR: δ = 0.05 (m, 12H), 0.54 (s, 3H), 0.85 (s, 9H), 0.89 (s, 9H), 1.13 (d, 3H), 1.22 (s, 3H), 1.23 (s, 3H), 1.00-2.20 (m, 15H), 2.30 (bd, 1H), 2.53 (dd, 1H), 2.86 (m, 1H), 3.27 (m, 1H), 3.45 (m, 1H), 3.55 (s, 1H), 3.83 (m, 1H), 4.21 (m, 1H), 4.52 (m, 1H), 4.93 (m, 1H), 4.98 (m, 1H), 5.79 (d, 1H, J = 11.4), 6.45 (d, 1H, J = 11.4).

Preparation 13: 1(S),3(R)-Bis-[tert-butyl(dimethylsilyl)oxy]-20(R)-(3-hydroxy-3-methyl-1-butoxy-9,10-secopregna-5(Z),7(E),10(19)-triene (Compound 21)

A solution under nitrogen of compound 7 (40 mg) in dichloromethane (4 ml) containing anthracene (20 mg) and triethylamine (50 μ l) in a Pyrex flask was irradiated with light from a high pressure ultra-violet lamp (type TQ150Z2 (Hanau) at 15°C for 30 minutes. The reaction mixture was filtered and concentrated in vacuo to give a residue. Purification by chromatography (silica gel, 30% ether in petroleum ether as eluent) gave 21.

NMR in agreement with structure.

Preparation 14: Compound 11 (R^3 = 4-(tetrahydro-4H-pyran-2-yloxy)-4-ethyl-1-hexyl)

To a solution of compound 3 (561 mg, 1.0 mmol) in dry tetrahydrofuran (10 ml) potassium tert-butoxide (0.4 g, 3.6

mmol), 18-Crown-6 (80 mg) and 2-(6-bromo-3-ethyl-3-hexyloxy)-tetrahydro-4H-pyran (Preparation 5b) (1.08 g, 3.68 mmol) were added. The mixture was stirred overnight and diluted with ethyl acetate (60 ml), then washed with water (3 x 10 ml) and saturated aqueous sodium chloride (10 ml), dried over MgSO_4 and concentrated in vacuo. The compound was then purified by chromatography (150 g silica gel, 10% ether in petroleum ether as eluant) to give the desired compound as a colourless oil.

10 NMR: δ = 0.05 (m, 12H), 0.55 (s, 3H), 0.82 (m, 6H), 0.86 (s, 9H), 0.89 (s, 9H), 1.07 (d, 3H, J = 6), 1.0-2.1 (m, 25 H), 2.03 (bt, 1H), 2.18 (bd, 1H), 2.30 (bd, 1H), 2.54 (dd, 1H), 2.87 (bd, 1H), 3.12 (m, 1H), 3.25 (m, 1H), 3.42 (m, 1H), 3.55 (m, 1H), 3.95 (m, 1H), 4.21 (m, 1H),
15 4.52 (m, 1H), 4.68 (m, 1H), 4.92 (bs, 1H), 4.98 (bs, 1H), 5.79 (d, 1H, J = 11), 6.46 (d, 1H, J = 11) ppm.

Preparation 15: Compound 10 (R^3 = 4-(tetrahydro-4H-pyran-2-yloxy)-4-ethyl-1-pentyl)

20 By following the procedure of Preparation 14 and substituting 2-(5-bromo-2-methyl-2-pentyloxy)-tetrahydro-4H-pyran for 2-(6-bromo-3-ethyl-3-hexyloxy)-tetrahydro-4H-pyran, the desired compound was obtained as a colourless oil.

25 NMR: δ = 0.05 (m, 12H), 0.55 (s, 3H), 0.86 (s, 9H), 0.89 (s, 9H), 1.07 (d, 3H, J = 6), 1.19 (s, 3H), 1.20 (s, 3H), 0.9-2.0 (m, 21 H), 2.03 (m, 1H), 2.16 (bd, 1H), 2.30 (bd, 1H), 2.55 (dd, 1H), 2.87 (bd, 1H), 3.15 (m, 1H), 3.25 (m, 1H), 3.43 (m, 1H), 3.55 (m, 1H), 3.93 (m, 1H), 4.21 (m,
30 1H), 4.52 (m, 1H), 4.71 (m, 1H), 4.93 (bs, 1H), 4.98 (bs, 1H), 5.80 (d, 1H, J = 11), 6.46 (d, 1H, J = 11) ppm.

Preparation 16: Compound 24 (R^3 = 4-(tetrahydro-4H-pyran-2-yloxy)-4-methyl-1-pentyl)

35 A solution of the compound 10 prepared in Preparation 15 (200 mg, 0.27 mmol), anthracene (200 mg, 1.1 mmol) and triethylamine (1 drop) in dichloromethane (15 ml) under nitrogen in a Pyrex flask was irradiated with light from a

high pressure ultraviolet lamp, type TQ150Z2 (Hanau) at about 10°C for 30 minutes. The reaction blend was filtered, concentrated in vacuo and purified by chromatography (30 g silica gel, 50% ether in petroleum ether as eluant) to give the desired compound as a colourless oil.

NMR: δ = 0.05 (m, 12H), 0.53 (s, 3H), 0.87 (m, 18H), 1.06 (d, 3H, J = 6), 1.18 (s, 3H), 1.20 (s, 3H), 1.0-1.9 (m, 21 H), 1.98 (bt, 1H), 2.16 (m, 2H), 2.43 (dd, 1H), 2.82 (bd, 1H), 3.18 (m, 1H), 3.24 (m, 1H), 3.43 (m, 1H), 3.53 (m, 1H), 3.93 (m, 1H), 4.18 (m, 1H), 4.36 (m, 1H), 4.70 (m, 1H), 4.85 (bs, 1H), 5.16 (bs, 1H), 5.99 (d, 1H, J = 11), 6.24 (d, 1H, J = 11) ppm.

Preparation 17: Compound 25 (R^3 = 4-(tetrahydro-4H-pyran-2-yloxy)-4-ethyl-1-hexyl)

By following the procedure of Preparation 16 and substituting the compound 11 prepared in Preparation 14 for the compound 10 prepared in Preparation 15, the desired compound was obtained as a colourless oil.

NMR: δ = 0.05 (m, 12H), 0.53 (s, 3H), 0.82 (m, 6H), 0.87 (s, 18H), 1.06 (d, 3H, J = 6), 1.0-1.9 (m, 25 H), 1.98 (bt, 1H), 2.19 (m, 2H), 2.44 (dd, 1H), 2.82 (bd, 1H), 3.12 (m, 1H), 3.25 (m, 1H), 3.43 (m, 1H), 3.55 (m, 1H), 3.93 (m, 1H), 4.18 (m, 1H), 4.36 (m, 1H), 4.69 (m, 1H), 4.85 (bs, 1H), 5.16 (bs, 1H), 5.99 (d, 1H, J = 11), 6.24 (d, 1H, J = 11) ppm.

Preparation 18: Compound 15 (R^3 = 5-trimethylsilyl oxy-5-methyl-1-hexyl)

To a solution of 3 (561 mg, 1.0 mmol) in dry tetrahydrofuran (10 ml) potassium tert-butoxide (0.65 g, 5.8 mmol), 18-Crown-6 (120 mg) and 6-bromo-2-methyl-2-trimethylsilyloxy-hexan (1.4 ml, 5.0 mmol) were added. The reaction mixture was stirred for 2 hours and worked-up (ether). The crude product was purified by chromatography (40 g silica gel, 2% ether in petroleum ether as eluant) to give a colourless oil, which crystallized from methanol.

M.p.: 75.5 - 77.5°C.

NMR: δ = 0.05-0.09 (m, 21H), 0.55 (s, 3H), 0.86 (s, 9H), 0.89 (s, 9H), 1.07 (d, 3H), 1.18 (s, 6H), 1.15-2.0 (m, 17H), 2.02 (t, 1H), 2.17 (d, 1H), 2.31 (d, 1H), 2.55 (dd, 1H), 2.85 (bd, 1H), 3.15 (m, 1H), 3.26 (m, 1H), 3.56 (m, 1H), 4.21 (m, 1H), 4.53 (m, 1H), 4.93 (bs, 1H), 4.99 (bs, 1H), 5.79 (d, 1H), 6.46 (d, 1H) ppm.

Preparation 19: Compound 16 (R^3 = 5-trimethylsilyl-oxy-5-ethyl-1-heptyl)

10 To a solution of compound 3 (561 mg, 1.0 mmol) in dry tetrahydrofuran (10 ml), potassium tert-butoxide 0.45 g, 4.0 mmol), 18-Crown-6 (80 mg) and 7-bromo-3-ethyl-3-trimethylsilyloxy-heptane (0.44 ml, 1.5 mol) were added. The reaction blend was stirred for 4 hours and worked-up (ethyl acetate). The crude product was purified by chromatography (100 g silica gel, 5% ether in petroleum ether as eluant) to give a colourless oil which crystallized from methanol.

M.p.: 70.5 - 72.5°C.

NMR: δ = 0.04-0.10 (m, 21H), 0.55 (s, 3H), 0.80 (dt, 6H), 0.86 (s, 9H), 0.89 (s, 9H), 1.07 (d, 3H), 1.43 (dq, 4H), 1.00-1.96 (m, 17H), 2.04 (bt, 1H), 2.17 (bd, 1H), 2.30 (bd, 1H), 2.55 (dd, 1H), 2.86 (bd, 1H), 3.15 (m, 1H), 3.26 (m, 1H), 3.58 (m, 1H), 4.21 (m, 1H), 4.52 (m, 1H), 4.93 (m, 1H), 4.98 (m, 1H), 5.80 (d, 1H, J = 11.3), 6.46 (d, 1H, J = 11.3) ppm.

Preparation 20: 1-(tert-butyl-dimethylsilyloxy)-6-chloro-hexane

To a solution of 6-chloro-hexan-1-ol (6.8 ml, 75.4 mmol) in dry dichloromethane (100 ml) tert-butyldimethylsilylchloride (12.5 g, 83 mmol) and imidazol (10.21 g, 150 mmol) were added, and the reaction mixture stirred overnight at room temperature. Work-up (dichloromethane) and distillation gave the title compound as an oil.

B.p.: 130-134°C/12 mBar.

NMR: δ = 0.03 (s, 6H), 0.88 (s, 9H), 1.27-1.60 (m, 6H), 1.77 (m, 2H), 3.52 (t, 2H), 3.59 (t, 2H) ppm.

Preparation 21: 1-(tert-butyldimethylsilyloxy)-6-iodo-hexane

A solution of sodium iodide (13.5 g, 90 mmol) and 1-(tert-butyldimethylsilyloxy)-6-chloro-hexane (Preparation 20) (8.35 g, 22 mmol) in acetone (70 ml) was refluxed overnight. The reaction mixture was cooled to room temperature and filtered. The filtrate was worked-up (hexane) to give the desired compound as yellow oil.

NMR: δ = 0.03 (s, 6H), 0.88 (s, 9H), 1.22-1.60 (m, 6H), 1.82 (m, 2H), 3.18 (t, 2H), 3.59 (t, 2H) ppm.

Preparation 22: Compound 17 (R^3 = 6-(tert-butyldimethylsilyloxy)-1-hexyl)

To a solution of compound 3 (516 mg, 0.9 mmol) in dry tetrahydrofuran (8 ml), potassium tert-butoxide (0.65 g, 5.8 mmol), 18-Crown-6 (100 mg) and 1-(tert-butyldimethylsilyloxy)-6-iodo-hexane (Preparation 21) (1.70 ml, 5 mmol) were added. The mixture was stirred overnight and worked-up (ether). The crude product was purified by chromatography (100 g silica gel, 30% toluene in petroleum ether as eluant) to give a colourless oil which crystallized from methanol.

M.p.: 84-87°C.

NMR: δ = 0.03 (s, 6H), 0.06 (m, 12H), 0.54 (s, 3H), 0.86 (s, 9H), 0.87 (s, 9H), 0.89 (s, 9H), 1.07 (d, 3H), 1.10-1.82 (m, 18H), 1.92 (m, 1H), 2.03 (bt, 1H), 2.14 (bd, 1H), 2.30 (bd, 1H), 2.52 (dd, 1H), 2.87 (m, 1H), 3.22 (m, 2H), 3.55 (m, 1H), 3.58 (t, 2H), 4.21 (m, 1H), 4.52 (m, 1H), 4.93 (m, 1H), 4.98 (m, 1H), 5.80 (d, 1H, J = 11.4), 6.46 (d, 1H, J = 11.4) ppm.

Preparation 23: Compound 30 (R^3 = 6-(tert-butyldimethylsilyloxy)-1-hexyl)

A solution of the compound 17 prepared in Preparation 22 (238 mg, 0.3 mmol), anthracene (150 mg, 0.8 mmol) and triethylamine (2 drops) in dichloromethane (12 ml) under nitrogen in a Pyrex flask was irradiated with light from a high pressure ultra-violet lamp, type TQ150Z2 (Hanau) at

15°C for 30 minutes. The reaction mixture was filtered, concentrated in vacuo and purified by chromatography (40 g silica gel, 10% ether in petroleum ether as eluant) to give the desired compound as a colourless oil.

5 NMR: δ = 0.03 (s, 6H), 0.04 (m, 6H), 0.05 (s, 6H), 0.53 (s, 3H), 0.86 (s, 9H), 0.87 (s, 9H), 0.88 (s, 9H), 1.06 (d, 3H), 1.00-2.30 (m, 22H), 2.44 (dd, 1H), 2.82 (bd, 1H), 3.20 (m, 2H), 3.55 (m, 1H), 3.58 (t, 2H), 4.18 (m, 1H), 4.36 (m, 1H), 4.86 (m, 1H), 5.16 (m, 1H), 5.99 (d, 1H, 10 J = 11.3), 6.24 (d, 1H, J = 11.3) ppm.

Preparation 24: Compound 29 (R^3 = 5-trimethylsilyloxy-5-ethyl-1-heptyl)

A solution of the compound 16 prepared in Preparation 15 19 (300 mg, 0.4 mmol), anthracene (300 mg, 1.7 mmol) and triethylamine (1 drop) in dichloromethane (15 ml) under nitrogen in a Pyrex flask was irradiated with light from a high pressure ultra-violet lamp, type TQ150Z2 (Hanau) at 15°C for 45 minutes. The reaction mixture was filtered, 20 concentrated in vacuo and purified by chromatography (15 g silica gel, 30% toluene in petroleum ether as eluant) to give the desired compound as a colourless oil.

NMR: δ = 0.05 (s, 6H), 0.06 (s, 6H), 0.08 (s, 9H), 0.54 (s, 3H), 0.80 (ds, 6H), 0.87 (s, 18H), 1.07 (d, 3H), 25 1.43 (bq, 4H), 1.00-2.25 (m, 20H), 2.45 (dd, 1H), 2.82 (bd, 1H), 3.15 (m, 1H), 3.24 (m, 1H), 3.57 (m, 1H), 4.18 (m, 1H), 4.35 (m, 1H), 4.86 (m, 1H), 5.16 (m, 1H), 5.99 (d, 1H, J = 11.3), 6.24 (d, 1H, J = 11.3) ppm.

30 Preparation 25: Compound 28 (R^3 = 5-trimethylsilyloxy-5-methyl-1-hexyl)

A solution of the compound 15 prepared in Preparation 18 (3.50 g, 4.7 mmol), anthracene (2.2 g, 12 mmol) and triethylamine (0.5 ml) in dichloromethane (175 ml) under 35 nitrogen in a Pyrex flask was irradiated with light from a high pressure ultra-violet lamp, type TQ150Z2 (Hanau) at 15°C for 2 hours. The reaction mixture was filtered, concentrated in vacuo and purified by chromatography (75 g

silica gel, 5% ether in petroleum ether as eluant) to give the desired compound as a colourless oil.

NMR: δ = 0.05-0.10 (m, 21H), 0.54 (s, 3H), 0.87 (s, 18H), 1.06 (d, 3H), 1.18 (s, 6H), 1.15-1.90 (m, 17H), 1.99 (t, 1H), 2.15 (m, 1H), 2.17 (m, 1H), 2.44 (dd, 1H), 2.81 (m, 1H), 3.20 (m, 2H), 3.56 (m, 1H), 4.18 (m, 1H), 4.36 (m, 1H), 4.86 (bd, 1H), 5.16 (bs, 1H), 5.98 (d, 1H), 6.23 (d, 1H) ppm.

10 Preparation 26: Compound 12 (R^3 = 4-trimethylsilyl-oxy-4-ethyl-1-hexyl)

A solution of potassium tert-butoxide (1.95 g, 17 mmol) in dry tetrahydrofuran (15 ml) was added dropwise via a syringe over 40 minutes to a solution of compound 3 (1.68 g, 3 mmol), 18-Crown-6 (600 mg) and 6-bromo-3-ethyl-3-trimethylsilyloxy-hexane (2.53 ml, 9 mmol) in dry tetrahydrofuran (20 ml) stirred under nitrogen. The resulting solution was stirred for 45 minutes and worked-up (hexane). The crude product was purified by chromatography (140 g silica gel, 30% toluene in petroleum ether as eluant) to give a colourless oil which crystallized from methanol.

M.p.: 52-57°C.

NMR: δ = 0.05-0.1 (m, 21H), 0.55 (s, 3H), 0.80 (dt, 6H), 0.86 (s, 9H), 0.89 (s, 9H), 1.07 (d, 3H), 1.10-2.05 (m, 20H), 2.18 (d, 1H), 2.30 (d, 1H), 2.54 (dd, 1H), 2.86 (bd, 1H), 3.12 (m, 1H), 3.25 (m, 1H), 3.55 (m, 1H), 4.21 (m, 1H), 4.52 (m, 1H), 4.93 (bs, 1H), 4.98 (bs, 1H); 5.79 (d, 1H), 6.46 (d, 1H) ppm.

30 Preparation 27: Compound 26 (R^3 = 4-trimethylsilyl-oxy)-4-ethyl-1-hexyl)

A solution of the compound 12 prepared in Preparation 26 (1.0 g, 1.3 mmol), anthracene (1.0 g, 5.6 mmol) and triethylamine (3 drops) in dichloromethane (70 ml) under nitrogen in a Pyrex flask was irradiated with light from a high pressure ultra-violet lamp, type TQ150Z2 (Hanau) at 15°C for 55 minutes. The reaction mixture was filtered, concentrated in vacuo, and purified by chromatography (35 g

silica gel, 2% ether in petrolum ether as eluant) to give the desired compound as a colourless oil.

NMR: δ = 0.05-0.10 (m, 21H), 0.54 (s, 3H), 0.80 (dt, 6H), 0.87 (s, 18H), 1.06 (d, 3H), 1.0-2.05 (m, 20H), 2.16 (d, 1H), 2.20 (m, 1H), 2.43 (dd, 1H), 2.81 (dd, 1H), 3.12 (m, 1H), 3.24 (m, 1H), 3.55 (m, 1H), 4.18 (m, 1H), 4.35 (m, 1H), 4.85 (bd, 1H), 5.16 (bs, 1H), 5.98 (d, 1H), 6.23 (d, 1H) ppm.

10 Preparation 28: Compound 18 (R^3 = 6-methyl-6-trimethylsilyloxy-1-heptyl)

A solution of 18-Crown-6 (264 mg, 1 mmol) in dry tetrahydrofuran (4 ml) was added dropwise via a syringe over 3 minutes to a mixture of compound 3 (561 mg, 1 mmol),
15 7-bromo-2-methyl-2-trimethylsilyloxy-heptane (1.5 ml, 4 mmol) and potassium hydride (0.6 ml, 20% suspension in oil) stirred under nitrogen. The resulting solution was stirred for 3 hours and worked-up (ether). The crude product was purified by chromatography (75 g silica gel, 5% ether in
20 petroleum ether as eluant) to give a colourless oil which crystallized from methanol.

NMR: δ = 0.06 (m, 12H), 0.08 (s, 9H), 0.54 (s, 3H), 0.86 (s, 9H), 0.89 (s, 9H), 1.07 (d, 3H), 1.18 (s, 6H), 1.00-1.83 (m, 18H), 1.90 (m, 1H), 2.03 (bt, 1H), 2.15 (bd, 1H), 2.31 (bd, 1H), 2.55 (dd, 1H), 2.87 (bd, 1H), 3.20 (m, 1H), 3.53 (m, 1H), 4.21 (m, 1H), 4.53 (m, 1H), 4.93 (m, 1H), 4.98 (m, 1H), 5.80 (d, 1H, J = 11.4), 4.46 (d, 1H, J = 11.4) ppm.

30 Preparation 29: Compound 31 (R^3 = 6-methyl-6-trimethylsilyloxy-1-heptyl)

A solution of the compound 18 prepared in Preparation 28 (400 mg, 0.152 mmol), anthracene (300 mg, 1.7 mmol) and triethylamine (3 drops) in dichloromethane (20 ml) under
35 nitrogen in a Pyrex flask was irradiated with light from a high pressure ultra-violet lamp, type TQ150Z2 (Hanau) at 15°C for 30 minutes. The reaction mixture was filtered, concentrated in vacuo and purified by chromatography (35 g

silica gel, 5% ether in petroleum ether as eluant) to give the desired compound as a colourless gum.

NMR: δ = 0.05 (m, 12H), 0.08 (s, 9H), 0.53 (s, 3H), 0.87 (s, 18H), 1.06 (d, 3H), 1.18 (s, 6H), 1.00-2.30 (m, 22H), 2.44 (dd, 1H), 2.81 (bd, 1H), 3.21 (m, 2H), 3.54 (m, 1H), 4.18 (m, 1H), 4.37 (m, 1H), 4.86 (m, 1H), 5.17 (m, 1H), 5.99 (d, 1H, J = 11.3), 6.24 (d, 1H, J = 11.3) ppm.

10 Preparation 30: 1-(tert-butyldimethylsilyloxy)-4-chloro-butane

To a solution of 4-chloro-butan-1-ol (10 ml, 100 mmol) in dry dichlormethane (100 ml) tert-butyldimethylsilylchloride (20.8 g, 120 mmol) and imidazol (13.61 g, 200 mmol) were added, and the reaction mixture was stirred 15 overnight at room temperature. Work-up (ethyl acetate) distillation gave the title compound as an oil.

B.p.: 89-92°C/12 mBar.

NMR: δ = 0.04 (s, 6H), 0.88 (s, 9H), 1.65 (m, 2H), 1.84 (m, 2H), 3.56 (t, 2H), 3.63 (t, 2H) ppm.

20

Preparation 31: Compound 9 (R^3 = 4-(tert-butyldimethylsilyloxy)-1-butyl)

A solution of 18-Crown-6 (264 mg, 1 mmol) in dry tetrahydrofuran (4 ml) was added dropwise via a syringe 25 over 2 minutes to a mixture of compound 3 (561 mg, 1 mmol) 4-chloro-1-tert-butyldimethylsilyloxy-butane (Preparation 30) (1.5 ml, 5 mmol) and potassium hydride (0.6 ml, 20% suspension in oil) stirred under nitrogen. The resulting solution was stirred for 3 1/2 hours and worked-up (ether). 30 The crude product was purified by chromatography (75 g silica gel, 5% ether in petroleum ether as eluant) to give a colourless gum which crystalized from methanol.

M.p.: 91-96°C.

35 NMR: δ = 0.03 (m, 6H), 0.06 (s, 12H), 0.54 (s, 3H), 0.86 (s, 9H), 0.88 (s, 9H), 0.89 (s, 9H), 1.06 (d, 3H), 1.00-1.83 (m, 14H), 1.92 (m, 1H), 2.03 (bt, 1H), 2.14 (bd, 1H), 2.30 (bd, 1H), 2.54 (dd, 1H), 2.86 (bd, 1H), 3.23 (m, 2H), 3.57 (dd, 1H), 3.61 (t, 2H), 4.21 (m, 1H), 4.53 (m,

1H), 4.93 (m, 1H), 4.98 (m, 1H), 5.79 (d, 1H, $J = 11.4$),
6.46 (d, 1H), $J = 11.4$) ppm.

5 Preparation 32: Compound 13 ($R^3 = 4$ -trimethylsilyl-
oxy-4-(1-propyl)-1-heptyl)

The compound was prepared according to the procedure described in Preparation 31, except that the 4-chloro-1-tert-butyl-dimethoxy-butane was substituted with 7-bromo-4-(1-propyl)-4-trimethylsilyloxy-heptane.

10 NMR: $\delta = 0.05$ (m, 12H), 0.07 (s, 9H), 0.55 (s, 3H),
0.86 (s, 9H), 0.87 (m, 6H), 0.89 (s, 9H), 1.07 (d, 3H),
1.00-1.85 (m, 22H), 1.91 (m, 1H), 2.03 (bt, 1H), 2.19 (bd,
1H), 2.30 (bd, 1H), 2.55 (dd, 1H), 2.87 (bd, 1H), 3.11 (m,
2H), 3.25 (dd, 1H), 3.55 (t, 1H), 4.22 (m, 1H), 4.53 (m,
15 1H), 4.93 (m, 1H), 4.93 (m, 1H), 4.98 (m, 1H), 5.80 (d, 1H,
 $J = 11.3$), 6.46 (d, 1H), $J = 11.3$) ppm.

20 Preparation 33: Compound 5 ($R^3 = 5$ -methyl-1-hexyl)

The compound was prepared according to the procedure described in Preparation 31, except that the 4-chloro-1-tert-butyl-dimethylsilyloxy-butane was substituted with 1-bromo-5-methyl-hexane.

M.p.: 79.5-81°C.

25 NMR: $\delta = 0.06$ (m, 12H), 0.55 (s, 3H), 0.85 (s, 6H),
0.86 (s, 9H), 0.89 (s, 9H), 1.07 (d, 3H), 1.00-1.85 (m,
17H), 1.91 (m, 1H), 2.03 (bt, 1H), 2.16 (bd, 1H), 2.31 (bd,
1H), 2.55 (dd, 1H), 2.87 (bd, 1H), 3.16 (m, 2H), 3.24 (m,
1H), 3.55 (m, 1H), 4.21 (m, 1H), 4.53 (m, 1H), 4.93 (m,
1H), 4.98 (m, 1H), 5.80 (d, 1H, $J = 11.4$), 6.46 (d, 1H), J
30 = 11.4) ppm.

35 Preparation 34: Compound 27 ($R^3 = 4$ -trimethylsilyl-
oxy-4-(1-propyl)-1-heptyl)

The compound was prepared according to the procedure described in Preparation 23, except that the compound 17 prepared in Preparation 22 was substituted with the compound 13 prepared in Preparation 32.

NMR: $\delta = 0.04$ (m, 6H), 0.05 (s, 6H), 0.07 (s, 9H),

0.54 (s, 3H), 0.80-0.93 (m, 24H), 1.06 (d, 3H), 1.00-2.07 (m, 24H), 2.19 (m, 2H), 2.45 (dd, 1H), 2.82 (bd, 1H), 3.12 (m, 1H), 3.24 (m, 2H), 3.55 (m, 1H), 4.18 (m, 1H), 4.36 (m, 1H), 4.86 (m, H), 5.17 (m, 1H), 5.99 (d, 1H, $J = 11.2$),
5 6.24 (d, 1H, $J = 11.2$) ppm.

Preparation 35: Compound 20 ($R^3 = 5\text{-methyl-1-hexyl}$)

The compound was prepared according to the procedure described in Preparation 23, except that compound 17
10 prepared in Preparation 22 was substituted with compound 5 prepared in Preparation 33.

NMR: $\delta = 0.05$ (s, 6H), 0.06 (s, 6H), 0.53 (s, 3H), 0.85 (d, 6H), 0.87 (s, 18H), 1.06 (d, 3H), 1.00-1.92 (m, 18H), 1.98 (bt, 1H), 2.18 (m, 2H), 2.44 (dd, 1H), 2.82 (bd, 1H), 3.18 (m, 2H), 3.55 (m, 1H), 4.17 (m, 1H), 4.36 (m, 1H), 4.86 (m, 1H), 5.16 (m, 1H), 5.99 (d, 1H, $J = 11.3$),
15 6.24 (d, 1H, $J = 11.3$) ppm.

Preparation 36: Compound 23 ($R^3 = 4\text{-tert-butyltrimethylsilyloxy-1-butyl}$)

The compound was prepared according to the procedure described in Preparation 23, except that compound 17 prepared in Preparation 22 was substituted with compound 9 prepared in Preparation 31.

25 NMR: $\delta = 0.05$ (m, 18H), 0.53 (s, 3H), 0.87 (m, 27H), 1.06 (d, 3H), 1.00-2.30 (m, 18H), 2.44 (dd, 1H), 2.82 (bd, 1H), 3.22 (m, 2H), 3.57 (m, 1H), 3.61 (t, 1H), 4.18 (m, 1H), 4.36 (m, 1H), 4.86 (m, 1H), 5.17 (m, 1H), 5.99 (d, 1H, $J = 11.3$), 6.24 (d, 1H, $J = 11.3$) ppm.

30

Preparation 37: Compound 8 ($R^3 = 2\text{-(1-hydroxycyclohexyl)ethyl}$)

Using the procedure of Preparation 11, but substituting 2-cyclohexylidene-1-bromoethane (2.5 g) for
35 the 3,3-dimethylallyl bromide the intermediate compound III ($R^3 = \text{CH}_2\text{-CH=C-(CH}_2\text{)}_4\text{-CH}_2$) was obtained. This compound (100 mg) was substituted for compound 4 in the procedure of Preparation 12 to give 8.

NMR in agreement with structure.

Preparation 38: Compound 22 ($R^3 = 2-(1\text{-hydroxycyclohexyl})\text{ethyl}$)

5 The compound was prepared using the procedure of Preparation 13, but substituting compound 8 for compound 7. NMR in agreement with structure.

10 Preparation 39: Compound 19 ($R^3 = 7\text{-methyl-7-trimethylsilyloxy-1-octyl}$)

The compound was prepared using the procedure of Preparation 28, but substituting 8-bromo-2-methyl-2-trimethylsilyloxy-octane for the 7-bromo 2-methyl-2-trimethylsilyloxy-heptane.

15 NMR in agreement with structure.

The 8-bromo-2-methyl-2-trimethylsilyloxy-octane used in the Preparation was prepared analogously to the method described for lower homologues in our PCT/DK89/00079.

B.p.: 92-95°C/0.1 mmHg.

20 NMR: $\delta = 0.09$ (s, 9H), 1.18 (s, 6H), 1.2-1.5 (m, 8H), 1.85 (m, 2H), and 3.40 (t, 2H).

Preparation 40: Compound 32 ($R^3 = (7\text{-methyl-7-trimethylsilyloxy-1-octyl})$)

25 The compound was prepared using the procedure of Preparation 29, but substituting compound 19 for compound 18.

NMR in agreement with structure.

30 Example 1: 1(S),3(R)-Dihydroxy-20(R)-(5'-hydroxy-5'-methyl-1'-hexyloxy)-9,10-seco-pregna-5(Z),7(E),10(19)-triene (Compound 109)

35 The compound 33 prepared in Preparation 4 (60 mg, 0.11 mmol) was dissolved in ethyl acetate (0.5 ml) and acetonitrile (5 ml) was added. A 5% solution of hydrofluoric acid in acetonitrile/water 8:1 (0.5 ml) was added, and the solution was stirred under nitrogen for 1

hour. Ethyl acetate (50 ml) was added, and the mixture was extracted with saturated aqueous sodium hydrogen carbonate (10 ml) and water (10 ml), dried and evaporated in vacuo. The residue was purified by chromatography (silica gel, ethyl acetate as eluant) to give 109.

NMR: δ = 0.56 (s, 3H), 1.07 (d, 3H), 1.20 (s, 6H), 1.10-2.05 (m, 24H), 2.15 (bd, 1H), 2.30 (dd, 1H), 2.60 (dd, 1H), 2.72 (m, 1H), 3.20 (m, 2H), 3.57 (m, 1H), 4.21 (m, 1H), 4.42 (m, 1H), 5.00 (bs, 1H), 5.32 (bs, 1H), 5.99 (d, 1H), 6.38 (d, 1H) ppm.

The same compound was obtained when compound 28 (Preparation 25) was used as starting material instead of compound 33.

Example 2: 1(S),3(R)-Dihydroxy-20(R)-(2'-hydroxy-3'-methyl-1'-butoxy)-9,10-seco-pregna-5(Z),7(E),10(19)-triene (Compound 101)

A solution of 1(S),3(R)-dihydroxy-20(R)-2'-hydroxy-3'-methyl-1'-butoxy)-9,10-secopregna-5(E),7(E),10(19)-triene (Compound 34) (80 mg) triethylamine (0.2 ml) and anthracene (50 mg) in dichloromethane (8 ml) was illuminated with a radiation from a high pressure ultra-violet lamp (type TQ 150Z; Hanau) for 1 hour. The solution was then filtered and concentrated and the product purified by chromatography (silica gel; ethyl acetate as eluant) to give the title compound. The approx. 1:1 mixture of 2'-epimers obtained from the sequence involving the racemic side chain building block gave the following NMR: δ = 0.54 and 0.56 (2s, 3H), 0.89 (d, 3H, J = 6.8), 0.96 (d, 3H, J = 6.8), 1.02-2.10 (m, 19H), 2.30 (m, 1H), 2.58 (m, 1H), 2.82 (m, 1H), 3.10-3.61 (m, 5H), 4.21 (m, 1H), 4.41 (m, 1H), 4.98 (m, 1H), 5.31 (m, 1H), 5.98 (d, 1H, J = 11.2), 6.36 (d, 1H, J = 11.2).

Example 3: 1(S),3(R)-Dihydroxy-20(R)-(3'-hydroxy-3'-methyl-1'-butoxy)-9,10-seco-
-pregna-5(Z),7(E),10(19)-triene
(Compound 102)

5 A solution of compound 21 (35 mg) in acetonitrile (4 ml) and 40% aqueous hydrofluoric acid (0.2 ml) was stirred at room temperature under nitrogen for 1 hour. Ethyl acetate was added and the mixture was extracted with saturated sodium hydrogen carbonate solution and then
10 brine. The ethyl acetate solution was dried and concentrated in vacuo to give a residue which was purified by chromatography (silica gel, ethyl acetate as eluent) to give the title compound.

 NMR: δ = 0.54 (s, 3H), 1.12 (d, 3H), 1.21 (s, 3H),
15 1.23 (s, 3H), 1.35-2.20 (m, 17H), 2.30 (dd, 1H), 2.57 (dd, 1H), 2.81 (m, 1H), 3.25 (m, 1H), 3.44 (m, 1H), 3.55 (s, 1H), 3.82 (m, 1H), 4.21 (m, 1H), 4.42 (m, 1H), 4.98 (m, 1H), 5.31 (m, 1H), 5.98 (d, 1H, J = 11.3), 6.37 (d, 1H, J = 11.3).

20

Example 4: 1(S),3(R)-Dihydroxy-20(R)-(4'-hydroxy-4'-methyl-1'-pentyloxy)-9,10-
-seco-pregna-5(Z),7(E),10(19)-triene
(Compound 105)

25 Compound 24 prepared in Preparation 16 (128 mg, 0.2 mmol) was dissolved in ethyl acetate (0.2 ml) and acetonitrile (4.4 ml) was added under vigorous stirring. A solution of 5% hydrofluoric acid in acetonitrile/water 8:1 (1.94 ml) was added, and the reaction mixture was stirred
30 under nitrogen at room temperature for 45 minutes. The reaction mixture was worked-up (ethyl acetate) and purified by chromatography (35 g silica gel, 80% ethyl acetate in petroleum ether as eluant) to give the desired compound as a colourless oil.

35 NMR: δ = 0.54 (s, 3H), 1.08 (d, 3H), 1.20 (s, 6H), 1.05-2.50 (m, 21H), 2.59 (dd, 1H), 2.81 (bd, 1H), 3.25 (m, 2H), 3.54 (m, 1H), 4.21 (m, 1H), 4.42 (m, 1H), 4.99 (m, 1H), 5.31 (m, 1H), 5.98 (d, 1H, J = 11.3), 6.37 (d, 1, J =

5

0

1

20

25

30

SUBSTITUTE SHEET

Example 7: 1(S),3(R)-Dihydroxy-20(R)-(6'-
-hydroxy-1'-hexyloxy)-9,10-seco-
-pregna-5(Z),7(E),10(19)-triene
(Compound 111)

5 Compound 30 prepared in Preparation 23 (233 mg, 0.3 mmol) was dissolved in ethyl acetate (0.6 ml) and acetonitrile (8 ml) was added under vigorous stirring. A solution of 5% hydrofluoric acid in acetonitrile/water 8:1 (4.0 ml) was added, and the reaction mixture was stirred under
10 nitrogen at room temperature for 90 minutes. The reaction mixture was worked-up (ethyl acetate) and purified by chromatography (40 g silica gel, 80% ethyl acetate in petroleum ether as eluant) to give the desired compound as a colourless gum.

15 NMR: δ = 0.55 (s, 3H), 1.07 (d, 3H), 1.00-2.22 (m, 24H), 2.31 (dd, 1H), 2.60 (dd, 1H), 2.83 (bd, 1H), 3.22 (m, 2H), 3.55 (m, 1H), 3.64 (t, 2H), 4.23 (m, 1H), 4.43 (m, 1H), 5.00 (m, 1H), 5.32 (m, 1H), 6.00 (d, 1H, J = 11.3), 6.39 (d, 1H, J = 11.3) ppm.

20

Example 8: 1(S),3(R)-Dihydroxy-20(R)-(6'-hydr-
-oxy-6'-methyl-1'-heptyloxy)-9,10-
-seco-pregna-5(Z),7(E),10(19)-triene
(Compound 112)

25 A solution of compound 31 prepared in Preparation 29 (300 mg) and tetrabutylammonium fluoride trihydrate (1.16 g) was stirred under nitrogen at 60°C for 60 minutes. After cooling, the reaction mixture was worked-up (ethyl acetate) and purified by chromatography (35 g silica gel, 80% ethyl
30 acetate in petroleum ether as eluant) to give the desired compound as a colourless gum.

 NMR: δ = 0.55 s, 3H), 1.07 (d, 3H), 1.20 (s, 6H), 1.00-2.22 (m, 24H), 2.30 (dd, 1H), 2.60 (dd, 1H), 2.84 (bd, 1H), 3.22 (m, 1H), 3.55 (m, 1H), 4.22 (m, 1H), 4.43 (m, 1H), 5.00 (m, 1H), 5.32 (m, 1H), 6.00 (d, 1H, 11.3), 6.39
35 (d, 1H, J = 11.3) ppm.

Example 9: 1(S),3(R)-Dihydroxy-20(R)-(4'-hydr-
-oxy-4'-(1"-propyl)-1'-heptyloxy)-
-9,10-seco-pregna-5(Z),7(E),10(19)-
-triene (Compound 107)

5 The compound was prepared according to the procedure described in Example 4, except that compound 24 prepared in Preparation 16 was substituted with compound 27 prepared in Preparation 34.

NMR: δ = 0.55 (s, 3H), 0.91 (t, 6H), 1.09 (d, 3H),
10 1.1-2.05 (m, 25H), 2.15 (bd, 1H), 2.32 (dd, 1H), 2.60 (bd, 1H), 2.82 (m, 1H), 3.22 (m, 2H), 3.56 (m, 1H), 4.23 (m, 1H), 4.43 (m, 1H), 5.00 (bs, 1H), 5.32 (bs, 1H), 5.99 (d, 1H), 6.48 (d, 1H) ppm.

15 Example 10: 1(S),3(R)-Dihydroxy-20(R)-(5'-meth-
-yl-1'-hexyloxy)-9,10-seco-pregna-
-5(Z),7(E),10(19)-triene
(Compound 108)

20 The compound was prepared according to the procedure described in Example 4, except that compound 24 prepared in Preparation 16 was substituted with compound 20 prepared in Preparation 35.

NMR: δ = 0.56 (s, 3H), 0.86 (d, 6H), 1.07 (d, 3H),
1.00-2.07 (m, 21H), 2.16 (bd, 1H), 2.31 (dd, 1H), 2.60 (bd,
25 1H), 2.82 (bd, 1H), 3.20 (m, 2H), 3.55 (m, 1H), 4.23 (m, 1H), 4.43 (m, 1H), 5.00 (bs, 1H), 5.32 (m, 1H), 6.00 (d, 1H, J = 11.3), 6.39 (d, 1H, J = 11.3) ppm.

30 Example 11: 1(S),3(R)-Dihydroxy-20(R)-(4'-
-hydroxy-1'-butyloxy)-9,10-seco-
-pregna-5(Z),7(E),10(19)-triene
(Compound 104)

35 The compound was prepared according to the procedure described in Example 4, except that compound 24 prepared in Preparation 16 was substituted with compound 23 prepared in Preparation 36.

NMR: δ = 0.56 (s, 3H), 1.10 (d, 3H), 1.00-2.20 (m, 19H), 2.32 (dd, 1H), 2.62 (m, 1H), 2.84 (bd, 1H), 3.30 (m,

The compound was prepared using the procedure of
10 Example 3, but substituting compound 22 for compound 21.
NMR in agreement with structure.

The compound was prepared using the procedure of Example 8, but substituting compound 32 for compound 31. NMR in agreement with structure.

30

<u>Example 15</u>	<u>Dermatological Cream Containing</u>
	<u>Compound 106</u>

In 1 g almond oil was dissolved 0.05 mg 106. To this solution was added 40 g of mineral oil and 20 g of self-emulsifying beeswax. The mixture was heated to liquify. After the addition of 40 ml hot water, the mixture was mixed well. The resulting cream contains approximately 0.5 µg of 106 per gram of cream.


WHAT WE CLAIM IS:

1. A compound of the formula I

5

10

15




in which formula R stands for an alkyl group containing
 20 from 4 to 12 carbon atoms optionally substituted with a
 hydroxy group; and derivatives of the compounds of formula
 I in which one or more hydroxy groups have been transformed
 into -O-acyl or -O-glycosyl or phosphate ester groups, such
 masked groups being hydrolyzable in vivo.

25

2. A compound according to claim 1, in which R is a
 group of formula II

30



II

where n is an integer from 1 to 7; R^1 and R^2 , which may be
 the same or different, stand for hydrogen, C_1 - C_5 -alkyl,
 35 C_3 - C_7 -cycloalkyl, or taken together with the carbon atom
 (starred in formula II) bearing the group X, R^1 and R^2 can
 form a C_3 - C_8 carbocyclic ring; and X stands for hydrogen or
 hydroxy.

3. A diastereoisomer of a compound according to claims 1 or 2, in pure form; or a mixture of diastereoisomers of a compound according to claim 1 or 2.
- 5 4. A compound according to claim 1, selected from the group consisting of
- 10 a) 1(S),3(R)-Dihydroxy-20(R)-(4'-hydroxy-4'-ethyl-1'-hexyloxy)-9,10-seco-pregna-5(Z),7(E),10(19)-triene
- b) 1(S),3(R)-Dihydroxy-20(R)-(6'-hydroxy-1'-hexyloxy)-9,10-seco-pregna-5(Z),7(E),10(19)-triene
- 15 c) 1(S),3(R)-Dihydroxy-20(R)-(5'-hydroxy-5'-ethyl-1'-heptyloxy)-9,10-seco-pregna-5(Z),7(E),10(19)-triene
- d) 1(S),3(R)-Dihydroxy-20(R)-(5'-hydroxy-5'-methyl-1'-hexyloxy)-9,10-seco-pregna-5(Z),7(E),10(19)-triene
- 20 e) 1(S),3(R)-Dihydroxy-20(R)-(5'-methyl-1'-hexyloxy)-9,10-seco-pregna-5(Z),7(E),10(19)-triene
- f) 1(S),3(R)-Dihydroxy-20(R)-(4'-hydroxy-4'-(1"-propyl)-1'-heptyloxy)-9,10-seco-pregna-5(Z),7(E),10(19)-triene
- 25 g) 1(S),3(R)-Dihydroxy-20(R)-(4'-hydroxy-4'-methyl-1'-pentyloxy)-9,10-seco-pregna-5(Z),7(E),10(19)-triene
- 30 h) 1(S),3(R)-Dihydroxy-20(R)-(3'-hydroxy-3'-methyl-1'-butyloxy)-9,10-seco-pregna-5(Z),7(E),10(19)-triene.
5. A method for producing a compound of formula I of
- 35 claim 1 using the following steps, in which
- a) 1(S),3(R)-bis-(tert-butyldimethylsilyloxy)-20(S)formyl-9,10-seco-pregna-5(E),7(E),10(19)-triene is oxidized

with oxygen in the presence of a basic catalyst, to form 1(S),3(R)-bis-(tert-butyldimethylsilyloxy)-9,10-seco-pregna-5(E),7(E),10(19)-triene-20-one;


- 5 b) 1(S),3(R)-bis-(tert-butyldimethylsilyloxy)-9,10-seco-pregna-5(E),7(E),10(19)-triene-20-one is reduced e.g. by sodium borohydride, to form 1(S),3(R)-bis-(tert-butyldimethylsilyloxy)-9,10-seco-pregna-5(E),7(E),10(19)-triene-20(R)-ol;

10

- c) 1(S),3(R)-bis-(tert-butyldimethylsilyloxy)-9,10-seco-pregna-5(E),7(E),10(19)-triene-20(R)-ol is alkylated under basic conditions with a side chain building block of formula $Z-R^3$, in which Z is a leaving group such as
15 halogen, p-toluenesulphonyloxy or methanesulphonyloxy, to form a compound of formula III

20

25



- 30 in which R^3 is R as defined above or an analogue thereof or optionally a radical which can be converted to R;

- d) a compound of the above formula III is subjected to a triplet-sensitized photoisomerisation, and, if necessary,
35 conversion of R^3 to R and deprotection, to form the desired compound of formula I of claim 1 or an analogue thereof.

6. Intermediate for the synthesis of compounds of for-

mula I and analogues thereof which is 1(S),3(R)-bis(tert-butyl dimethylsilyloxy)-9,10-seco-pregna-5(E),7(E),10(19)-triene-20(R)-ol.

- 5 7. A pharmaceutical composition containing an effective amount of one or more of the compounds of claim 1, together with pharmaceutically acceptable, non-toxic carriers and/or auxiliary agents.
- 10 8. A pharmaceutical composition according to claim 7 in dosage unit form.
- 15 9. A dosage unit according to claim 8 containing from 0.05 - 50 µg, preferably from 0.1 - 25 µg of a compound of formula I.
- 20 10. A method for the treatment and prophylaxis of autoimmune diseases, including diabetes mellitus, hypertension, inflammatory diseases such as rheumatoid arthritis and asthma as well as diseases characterized by abnormal cell differentiation and/or cell proliferation, and/or imbalance in the immune system.
- 25 11. A method according to claim 10 for the treatment or prophylaxis of cancer.
12. A method according to claim 10 for the treatment of psoriasis.

INTERNATIONAL SEARCH REPORT.

International Application No PCT/DK 90/00036

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶ According to International Patent Classification (IPC) or to both National Classification and IPC IPC5: C 07 C 401/00, A 61 K 31/59														
II. FIELDS SEARCHED <div style="text-align: center; border: 1px solid black; padding: 2px;">Minimum Documentation Searched⁷</div> <table style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 20%; border: 1px solid black; padding: 2px;">Classification System</th> <th style="border: 1px solid black; padding: 2px;">Classification Symbols</th> </tr> <tr> <td style="border: 1px solid black; padding: 5px; vertical-align: top;">IPC5</td> <td style="border: 1px solid black; padding: 5px; vertical-align: top;">C 07 C; A 61 K</td> </tr> </table> <div style="text-align: center; border: 1px solid black; padding: 2px;">Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in Fields Searched⁸</div> <p style="padding: 5px;">SE,DK,FI,NO classes as above</p>			Classification System	Classification Symbols	IPC5	C 07 C; A 61 K								
Classification System	Classification Symbols													
IPC5	C 07 C; A 61 K													
III. DOCUMENTS CONSIDERED TO BE RELEVANT⁹ <table style="width: 100%; border-collapse: collapse;"> <tr> <th style="width: 10%; border: 1px solid black; padding: 2px;">Category¹</th> <th style="width: 70%; border: 1px solid black; padding: 2px;">Citation of Document,¹¹ with indication, where appropriate, of the relevant passages¹²</th> <th style="width: 20%; border: 1px solid black; padding: 2px;">Relevant to Claim No.¹³</th> </tr> <tr> <td style="border: 1px solid black; padding: 5px; vertical-align: top;">A</td> <td style="border: 1px solid black; padding: 5px; vertical-align: top;">EP, A2, 0184112 (CHUGAI SEIYAKU KABUSHIKI KAISHA) 11 June 1986, see the whole document --</td> <td style="border: 1px solid black; padding: 5px; vertical-align: top;">1-10</td> </tr> <tr> <td style="border: 1px solid black; padding: 5px; vertical-align: top;">A</td> <td style="border: 1px solid black; padding: 5px; vertical-align: top;">Chem. Pharm. Bull., Vol. 34, No. 10, 1986 Eigoro Murayama et al.: "Synthetic studies of vitamin D3 analogues. VIII. Synthesis of 22-oxavitamin D3 analogues. ", see page 4410 - page 4413 --</td> <td style="border: 1px solid black; padding: 5px; vertical-align: top;">1-10</td> </tr> <tr> <td style="border: 1px solid black; padding: 5px; vertical-align: top;">A</td> <td style="border: 1px solid black; padding: 5px; vertical-align: top;">Chemical Abstracts, volume 108, no. 11, 14 March 1988, (Columbus, Ohio, US), Abe Junko et al. : "Synthetic analogs of vitamin D3 with an oxygen atom in the side chain skeleton. ", see, abstract 88191v, & FEBS Lett. 1987, 226(1), 58- 62 --</td> <td style="border: 1px solid black; padding: 5px; vertical-align: top;">1-10</td> </tr> </table>			Category ¹	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³	A	EP, A2, 0184112 (CHUGAI SEIYAKU KABUSHIKI KAISHA) 11 June 1986, see the whole document --	1-10	A	Chem. Pharm. Bull., Vol. 34, No. 10, 1986 Eigoro Murayama et al.: "Synthetic studies of vitamin D3 analogues. VIII. Synthesis of 22-oxavitamin D3 analogues. ", see page 4410 - page 4413 --	1-10	A	Chemical Abstracts, volume 108, no. 11, 14 March 1988, (Columbus, Ohio, US), Abe Junko et al. : "Synthetic analogs of vitamin D3 with an oxygen atom in the side chain skeleton. ", see, abstract 88191v, & FEBS Lett. 1987, 226(1), 58- 62 --	1-10
Category ¹	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³												
A	EP, A2, 0184112 (CHUGAI SEIYAKU KABUSHIKI KAISHA) 11 June 1986, see the whole document --	1-10												
A	Chem. Pharm. Bull., Vol. 34, No. 10, 1986 Eigoro Murayama et al.: "Synthetic studies of vitamin D3 analogues. VIII. Synthesis of 22-oxavitamin D3 analogues. ", see page 4410 - page 4413 --	1-10												
A	Chemical Abstracts, volume 108, no. 11, 14 March 1988, (Columbus, Ohio, US), Abe Junko et al. : "Synthetic analogs of vitamin D3 with an oxygen atom in the side chain skeleton. ", see, abstract 88191v, & FEBS Lett. 1987, 226(1), 58- 62 --	1-10												
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>* Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance, the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance, the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div>														
IV. CERTIFICATION <table style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 50%; border: 1px solid black; padding: 5px;"> Date of the Actual Completion of the International Search 16th May 1990 </td> <td style="width: 50%; border: 1px solid black; padding: 5px;"> Date of Mailing of this International Search Report 1990 -05- 24 </td> </tr> <tr> <td style="border: 1px solid black; padding: 5px;"> International Searching Authority SWEDISH PATENT OFFICE </td> <td style="border: 1px solid black; padding: 5px;"> Signature of Authorized Officer Anna Hedberg <i>Anna Hedberg</i> </td> </tr> </table>			Date of the Actual Completion of the International Search 16th May 1990	Date of Mailing of this International Search Report 1990 -05- 24	International Searching Authority SWEDISH PATENT OFFICE	Signature of Authorized Officer Anna Hedberg <i>Anna Hedberg</i>								
Date of the Actual Completion of the International Search 16th May 1990	Date of Mailing of this International Search Report 1990 -05- 24													
International Searching Authority SWEDISH PATENT OFFICE	Signature of Authorized Officer Anna Hedberg <i>Anna Hedberg</i>													

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)

Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
A	<p>Chemical Abstracts, volume 110, no. 2, 9 January 1989, (Columbus, Ohio, US), Miyamoto Katsuhito et al. : "Antitumor pharmaceuticals containing 9,10-seco-5,7,10(19)-pregnatriene derivatives. ", see, abstract 13599s, & Jpn. Kokai Tokkyo Koho JP 63107930, 12 May 1988</p> <p>---</p> <p>-----</p>	1-10

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

V. ☒ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE ¹

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☒ Claim numbers 10-12 because they relate to subject matter not required to be searched by this Authority, namely:

See PCT Rule 39.1(iv): Methods for treatment of the human or animal body by surgery or therapy as well as diagnostic methods.

2. ☐ Claim numbers because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claim numbers because they are dependent claims and are not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☐ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING ²

This international Searching Authority found multiple inventions in this international application as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.
2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:
3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:
4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the international Searching Authority did not invite payment of any additional fee.

Remark on Protest

- ☐ The additional search fees were accompanied by applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

ANNEX TO THE INTERNATIONAL SEARCH REPORT
ON INTERNATIONAL PATENT APPLICATION NO.PCT/DK 90/00036

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.
The members are as contained in the Swedish Patent Office EDP file on 90-05-07
The Swedish Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A2- 0184112	86-06-11	JP-A- 61267548	86-11-27
		JP-A- 61267550	86-11-27